

Harnessing Microbial Metabolism for Sustainable Environmental Remediation

Rekha Rani¹; Dr. Neeraj Choudhary²

^{1,2}Department of Food and Biotechnology Faculty of Education and Methodology
Jayoti Vidyapeetham Women's University Jaipur Rajasthan

Publication Date: 2026/10/07

Abstract: Environmental contamination caused by industrialization, urbanization, intensive agriculture, petroleum exploitation, mining, pharmaceutical production, and improper waste disposal has created persistent challenges for ecosystem stability and human well-being. Conventional remediation technologies, including chemical oxidation, thermal treatment, excavation, incineration, and physicochemical separation, can be effective but may require substantial energy, chemical inputs, infrastructure, and secondary waste management. Microbial bioremediation provides an alternative approach by exploiting the metabolic versatility of bacteria, fungi, archaea, algae, and their associated enzymes to transform, degrade, immobilize, or recover environmental contaminants. This review examines how microbial metabolism can be harnessed for sustainable remediation of organic and inorganic pollutants. Particular attention is given to hydrocarbon degradation, pesticide and pharmaceutical transformation, dye decolorization, heavy-metal immobilization, plastic degradation, and treatment of complex industrial wastewater. Microbial remediation mechanisms include aerobic oxidation, anaerobic reduction, hydrolysis, dehalogenation, biosorption, bioaccumulation, biomineralization, cometabolism, and extracellular enzymatic transformation. The paper further examines bioaugmentation, biostimulation, microbial consortia, enzyme immobilization, metabolic engineering, omics-based approaches, phytomicrobial remediation, and microbial electrochemical technologies. Recent research indicates that microbial electrochemical systems can couple pollutant transformation with energy recovery and resource recovery, while advances in metagenomics, synthetic biology, artificial intelligence, biosensors, and engineered enzymes are expanding the potential of microbial remediation. Nevertheless, challenges remain concerning pollutant bioavailability, environmental variability, microbial survival, incomplete degradation, toxic intermediates, scale-up, ecological risks, and regulatory acceptance. Future remediation strategies should therefore move toward site-specific, mechanism-based systems that integrate microbial ecology, metabolic engineering, real-time monitoring, and resource recovery.

Keywords: Microbial Metabolism, Bioremediation, Biodegradation, Microbial Enzymes, Environmental Pollution, Bioaugmentation, Biostimulation, Microbial Consortia.

How to Cite: Rekha Rani; Dr. Neeraj Choudhary (2026) Harnessing Microbial Metabolism for Sustainable Environmental Remediation. *International Journal of Innovative Science and Research Technology*, 11(9), 3089-3096.
<https://doi.org/10.38124/ijisrt/26sep1362>

I. INTRODUCTION

Environmental pollution has become a multidimensional challenge involving soil, water, sediments, air, and biological systems. Industrial activities, petroleum extraction, agricultural intensification, mining, wastewater discharge, pharmaceutical manufacturing, plastics production, and inappropriate disposal of solid and hazardous wastes introduce diverse contaminants into ecosystems. These contaminants include petroleum hydrocarbons, polycyclic aromatic hydrocarbons (PAHs), pesticides, dyes, pharmaceuticals, phenolic compounds, chlorinated chemicals, heavy metals, metalloids, plastics, and emerging contaminants. Many of these substances are persistent because of their chemical stability, low bioavailability, toxicity, or resistance to conventional biological treatment.

Environmental remediation traditionally relies on physical and chemical technologies such as excavation, adsorption, coagulation, precipitation, oxidation, thermal treatment, membrane separation, and incineration. Although these methods can provide rapid contaminant removal, they may involve considerable energy consumption, chemical requirements, infrastructure, and secondary waste generation. These limitations have increased interest in biological approaches that use naturally occurring metabolic processes to reduce contaminant concentrations and toxicity.

Bioremediation exploits the capacity of microorganisms to use environmental chemicals as carbon sources, energy sources, electron donors, electron acceptors, or cometabolic substrates. Microorganisms can transform contaminants through intracellular metabolic pathways or extracellular enzymatic reactions. Enzymes such as

oxygenases, oxidoreductases, hydrolases, dehalogenases, laccases, and peroxidases participate in the transformation of numerous organic pollutants. Microbial systems can also immobilize or transform metals through biosorption, bioaccumulation, precipitation, redox transformation, and biomineralization rather than chemically degrading the metals themselves. (ScienceDirect)

The significance of microbial metabolism in environmental remediation arises from its diversity and adaptability. Microorganisms occupy almost every environmental niche and possess extensive metabolic capabilities. Microbial communities can cooperate through syntrophy, cross-feeding, cometabolism, extracellular enzyme production, and exchange of metabolic intermediates. In petroleum-contaminated environments, for example, mixed microbial communities may exhibit complementary substrate specificities that allow simultaneous transformation of different hydrocarbon fractions. (ScienceDirect)

Recent developments have extended microbial remediation beyond conventional biodegradation. Metagenomics and other omics approaches can reveal the metabolic potential of complex microbial communities, while synthetic biology and metabolic engineering can modify microorganisms or enzymes for specific transformation pathways. Microbial electrochemical technologies additionally exploit microbial electron-transfer processes to facilitate pollutant degradation while potentially generating electrical energy or recovering resources. (DOI)

Therefore, the central objective of this review is to examine how microbial metabolism can be deliberately harnessed for sustainable environmental remediation, with emphasis on mechanisms, pollutant-specific applications, technological strategies, limitations, and future directions.

II. OMICS AND SYSTEMS BIOLOGY FOR METABOLIC REMEDIATION

Omics technologies and systems biology have emerged as powerful approaches for understanding the complex microbial processes involved in environmental remediation. Conventional bioremediation research has frequently concentrated on individual microorganisms, isolated enzymes, or specific degradation pathways. Although these approaches have provided important information about pollutant transformation, they do not fully represent the complexity of natural microbial communities. Contaminated soils, sediments, groundwater, and wastewater contain diverse populations of bacteria, fungi, archaea, algae, and other microorganisms that interact with one another and respond dynamically to environmental conditions. Their metabolic activities are influenced by pollutant concentration, nutrient availability, temperature, pH, oxygen conditions, moisture, electron donors and acceptors, and interactions with other organisms. Omics and systems biology provide integrated approaches for investigating these complex relationships and identifying the molecular mechanisms that control contaminant transformation.

Consequently, their application is becoming increasingly important for developing targeted, efficient, and sustainable microbial remediation technologies.

The term omics encompasses a collection of high-throughput molecular approaches that investigate biological systems at different levels of organization. Important approaches for environmental remediation include genomics, metagenomics, transcriptomics, metatranscriptomics, proteomics, metaproteomics, metabolomics, and fluxomics. Genomics examines the complete genetic information of an individual microorganism and can identify genes associated with pollutant degradation, stress resistance, transport, energy metabolism, and detoxification. Metagenomics extends this analysis to entire microbial communities by examining DNA directly extracted from environmental samples. This is particularly valuable because many environmental microorganisms cannot be readily cultivated using conventional laboratory techniques. Metagenomic investigations can therefore reveal previously unknown microorganisms and functional genes associated with the degradation of petroleum hydrocarbons, pesticides, pharmaceuticals, aromatic compounds, dyes, plastics, and other environmental contaminants. By comparing microbial communities from contaminated and uncontaminated environments, researchers can identify organisms and functional genes that become enriched in response to particular pollutants.

Metagenomics is particularly useful for determining the metabolic potential of environmental microbial communities. For example, microorganisms inhabiting petroleum-contaminated soils may contain genes encoding alkane hydroxylases, monooxygenases, dioxygenases, dehydrogenases, and other enzymes involved in hydrocarbon transformation. Similarly, pesticide-contaminated environments may contain microorganisms possessing genes responsible for hydrolysis, oxidation, reduction, dehalogenation, and other reactions involved in pesticide degradation. The identification of these genes provides information about what microorganisms are potentially capable of doing. However, the presence of a particular gene does not necessarily indicate that the corresponding metabolic pathway is active. Environmental microorganisms can possess extensive genetic potential while expressing only a fraction of their genes under specific environmental conditions. This limitation highlights the importance of combining metagenomics with transcriptomics and other omics approaches.

Metatranscriptomics provides information about genes that are actively expressed within a microbial community. It involves the analysis of RNA transcripts and therefore provides a closer indication of microbial responses to environmental contaminants. For example, a contaminated soil may contain several microorganisms with genes associated with hydrocarbon degradation, but only certain microorganisms may actively express these genes when oxygen, nutrients, or other environmental conditions are favorable. Metatranscriptomic analysis can identify which metabolic pathways are activated during different stages of

remediation and can reveal how microorganisms respond to changing pollutant concentrations. This information can help researchers determine why a microbial population performs effectively under one set of conditions but becomes less active under another. It can also assist in optimizing biostimulation strategies by identifying the nutrients, electron donors, electron acceptors, or environmental conditions required to activate desirable metabolic pathways.

Proteomics adds another important level of information by examining proteins produced by microorganisms. Since enzymes and other proteins perform most cellular functions, proteomic analysis provides evidence of the biochemical processes actually occurring within microbial cells. In environmental remediation, proteomics can identify oxygenases, oxidoreductases, hydrolases, laccases, peroxidases, dehalogenases, transport proteins, stress-response proteins, and other proteins associated with pollutant transformation and microbial adaptation. Metaproteomics extends this approach to microbial communities and can determine how different members contribute to overall environmental processes. This is particularly important when pollutant degradation involves several successive reactions. One microorganism may transform a complex contaminant into an intermediate, while another microorganism may use that intermediate as a substrate and convert it into another compound. Protein-level analysis can therefore help identify the organisms and enzymes responsible for different stages of the remediation process.

Metabolomics provides complementary information by examining small molecules and metabolic products present within an environmental or biological system. Its importance in remediation lies in its ability to identify the products generated during pollutant transformation. A reduction in the concentration of a parent pollutant does not necessarily indicate complete detoxification because microorganisms may transform the contaminant into intermediate compounds that remain toxic or environmentally persistent. Metabolomic analysis can identify these intermediates and help determine whether degradation proceeds toward detoxification and mineralization. For example, the microbial transformation of an aromatic compound may involve several intermediate metabolites before the carbon is incorporated into central metabolic pathways. By monitoring these metabolites, researchers can reconstruct degradation pathways and determine whether particular biochemical steps are limiting the overall process.

The integration of genomics, transcriptomics, proteomics, and metabolomics provides a comprehensive multi-omics approach to environmental remediation. Each omics technology provides information at a different biological level, and their combination can establish stronger relationships between microbial community composition, genetic potential, gene expression, enzyme production, and metabolic activity. Metagenomics can reveal which degradation genes are present, metatranscriptomics can determine whether these genes are expressed, proteomics can identify the corresponding enzymes, and metabolomics can

reveal the products generated through their activity. Such integrated analysis provides a more accurate representation of microbial metabolism than any individual approach. It can also help identify metabolic bottlenecks, determine why particular degradation pathways are inefficient, and reveal environmental factors that control microbial activity.

Systems biology extends the application of omics by treating microorganisms and microbial communities as interconnected biological networks rather than isolated components. It integrates information about genes, proteins, metabolites, metabolic pathways, microbial interactions, and environmental variables to understand how biological systems function as a whole. In environmental remediation, systems biology can be used to construct metabolic networks showing how pollutants enter microbial pathways, how they are transformed into intermediate compounds, and how different microorganisms contribute to subsequent stages of degradation. Such network-based approaches are particularly important for understanding microbial consortia because complete pollutant degradation often requires cooperation among multiple microbial species. One microorganism may initiate the transformation of a pollutant, while another may consume the resulting intermediate, thereby preventing accumulation of potentially toxic products. These metabolic interactions can support more complete degradation than the activity of individual microorganisms.

Omics and systems biology can also contribute to the development of effective microbial consortia for bioremediation. Traditional bioaugmentation generally involves selecting individual microorganisms based on their ability to degrade a particular contaminant. However, individual strains may perform poorly under complex environmental conditions because they lack complementary metabolic functions or cannot compete successfully with native microbial populations. Systems-level analysis can identify microorganisms with complementary metabolic capabilities and reveal relationships such as syntrophy, cross-feeding, cooperation, and competition. Researchers can therefore design microbial communities in which different members perform different stages of a degradation pathway. Such designed consortia may be particularly valuable for treating petroleum hydrocarbons, mixed industrial contaminants, pharmaceutical residues, pesticides, and complex wastewater containing multiple pollutants.

The information obtained through omics approaches can further support metabolic engineering and synthetic biology. Once efficient pollutant-degrading genes and enzymes have been identified, researchers can investigate ways of enhancing their expression or transferring desirable metabolic functions into suitable microbial hosts. Systems biology can identify metabolic limitations and determine which steps restrict the efficiency of a degradation pathway. Metabolic engineering can then be used to modify microbial pathways to improve pollutant utilization, increase tolerance to toxic compounds, enhance enzyme production, or reduce the accumulation of harmful intermediates. Engineered enzymes may also be developed for controlled applications in treatment reactors or immobilized systems. Nevertheless, the

environmental application of genetically modified microorganisms requires careful consideration of ecological interactions, genetic stability, horizontal gene transfer, containment, and regulatory requirements.

The integration of omics with microbial electrochemical technologies represents another promising direction. Microbial electrochemical systems depend on microorganisms capable of transferring electrons to or from electrodes. Omics approaches can help identify electroactive microorganisms, characterize genes and proteins associated with extracellular electron transfer, and determine how microbial communities respond to electrode conditions. Systems-level analysis can subsequently be used to optimize electron flow toward pollutant degradation rather than competing metabolic pathways. This could improve the efficiency of bioelectrochemical systems for treating complex contaminants while potentially enabling simultaneous energy or resource recovery.

Artificial intelligence and machine-learning technologies are also expected to strengthen omics-based environmental remediation. Large datasets generated through metagenomics, metatranscriptomics, proteomics, metabolomics, environmental monitoring, and chemical analysis can be integrated using computational models. These models can identify relationships between environmental conditions, microbial community structure, metabolic activity, and pollutant removal. Predictive approaches may eventually help determine which microbial populations are most suitable for a particular contaminated site and which environmental conditions should be modified to stimulate their activity. Integration with real-time biosensors could allow treatment systems to continuously monitor pollutant concentrations and microbial responses and automatically adjust oxygen, nutrients, electron donors, pH, or other operational parameters.

Despite their considerable potential, omics and systems biology face several challenges. Environmental samples are highly complex, and separating signals from different microorganisms can be difficult. Large omics datasets require advanced computational resources, reliable databases, and sophisticated statistical and bioinformatic approaches. Furthermore, detecting a degradation gene or transcript does not necessarily prove that the corresponding pollutant is being completely degraded. Transformation products, enzyme activity, metabolic flux, and toxicity must also be evaluated. Standardization of sampling, sequencing, data analysis, and validation remains important for comparing results across different studies. The cost and technical complexity of multi-omics investigations may also restrict their application in routine field remediation.

III. MAJOR FACTORS INFLUENCING THE EFFICIENCY OF MICROBIAL REMEDIATION

The effectiveness of microbial remediation is governed by a complex interaction of biological, physicochemical, and environmental factors. Microorganisms require suitable

conditions for growth, reproduction, enzyme production, and metabolic activity, and any factor that limits these processes can reduce the efficiency of contaminant transformation. The nature and concentration of the pollutant are particularly important because different microorganisms possess different metabolic capabilities and tolerance levels. Highly toxic contaminants may inhibit microbial growth, whereas low concentrations may provide insufficient stimulation of specialized degraders. The chemical structure, solubility, hydrophobicity, molecular weight, and persistence of a pollutant also determine its susceptibility to microbial degradation. Pollutant bioavailability is equally important because contaminants strongly bound to soil particles or organic matter may remain inaccessible to microorganisms even when their total concentration is high. Consequently, successful remediation requires consideration of both contaminant concentration and the fraction that is biologically accessible.

pH is another major factor affecting microbial remediation because microbial growth and enzyme activity are strongly influenced by acidity and alkalinity. Most environmental microorganisms have relatively defined pH ranges within which their metabolic processes operate efficiently. Changes in pH can also alter the chemical form, solubility, adsorption, and bioavailability of pollutants. Extreme pH conditions may inhibit microbial growth or alter enzyme structure, thereby reducing degradation rates. Similarly, temperature influences microbial growth, membrane activity, enzyme kinetics, substrate availability, and overall metabolic rates. Microbial communities adapted to cold environments may possess enzymes that remain active at lower temperatures, whereas microorganisms from warmer environments may perform better under elevated temperatures. Therefore, remediation strategies should consider the natural temperature conditions of the contaminated site rather than relying exclusively on laboratory conditions.

The availability of essential nutrients such as nitrogen, phosphorus, sulfur, carbon, and trace elements can substantially influence microbial activity. In many contaminated environments, microorganisms may have sufficient carbon from pollutants but insufficient nitrogen or phosphorus for cellular growth. Biostimulation can address these limitations by supplying appropriate nutrients and thereby increasing the activity of indigenous microbial populations. However, excessive nutrient addition can cause undesirable effects, including eutrophication, changes in microbial community structure, and secondary environmental contamination. Nutrient supplementation should therefore be carefully controlled according to site-specific requirements. Oxygen availability is equally important because aerobic microorganisms require oxygen for many oxidative degradation pathways, whereas anaerobic microorganisms depend on alternative electron acceptors such as nitrate, sulfate, iron, or carbon dioxide. Maintaining suitable redox conditions can therefore determine whether particular microbial metabolic pathways remain active.

Moisture content is particularly important in soil bioremediation. Water is necessary for microbial metabolism, nutrient transport, enzyme activity, and movement of soluble contaminants. Excessive moisture, however, can restrict oxygen diffusion and promote anaerobic conditions, while insufficient moisture can inhibit microbial growth and reduce pollutant mobility. Soil texture, porosity, organic matter, salinity, and permeability can further influence the movement of microorganisms and contaminants. These factors demonstrate why a remediation approach that performs successfully in a laboratory reactor may not produce identical results under field conditions. Environmental heterogeneity can create localized zones with substantially different physicochemical and biological characteristics.

The composition and diversity of the microbial community are also fundamental determinants of remediation performance. Indigenous microorganisms may already possess specialized pathways for transforming contaminants because long-term exposure can select for adapted populations. Microbial diversity can provide functional redundancy, allowing the community to maintain degradation activity even when environmental conditions change. Conversely, low microbial diversity or the absence of appropriate metabolic functions can limit remediation. Bioaugmentation may therefore be considered when indigenous populations lack the required metabolic capabilities. However, introduced microorganisms must compete with established microbial communities and survive predation, environmental stress, and fluctuating conditions. Their long-term establishment cannot always be guaranteed.

Interactions among microorganisms can either enhance or inhibit remediation. Cooperation, syntrophy, cross-feeding, and metabolic complementarity may enable complete degradation of complex contaminants. One microorganism may initiate pollutant transformation while another consumes the resulting intermediate. Competition for nutrients or electron donors, predation, antagonistic interactions, and changes in community structure can have the opposite effect. Understanding microbial interactions is therefore essential for designing effective consortia and improving the predictability of biological remediation.

IV. EMERGING TECHNOLOGIES AND FUTURE DIRECTIONS IN MICROBIAL REMEDIATION

Rapid advances in biotechnology, molecular biology, computational science, and environmental engineering are creating new opportunities for improving microbial remediation. Conventional bioremediation often depends on naturally occurring microorganisms and environmental manipulation, but emerging approaches increasingly seek to identify, predict, and control microbial metabolic processes at molecular and community levels. The integration of omics technologies, synthetic biology, metabolic engineering, artificial intelligence, biosensors, microbial electrochemical systems, and advanced reactor technologies is creating a transition toward more precise and adaptive remediation. These developments may allow treatment systems to be

designed according to the specific contaminants, microbial communities, and environmental conditions present at individual sites.

Metagenomics, metatranscriptomics, proteomics, metabolomics, and other omics technologies are expected to play an increasingly important role in identifying microbial degradation pathways. These approaches can reveal previously unknown microorganisms, functional genes, enzymes, metabolic intermediates, and microbial interactions associated with pollutant transformation. Integrating multiple molecular datasets can help distinguish between microorganisms that merely possess degradation genes and those that actively contribute to contaminant removal. This information can improve the selection of microorganisms for bioaugmentation and identify environmental conditions that should be modified through biostimulation.

Metabolic engineering and synthetic biology represent another important area of development. Microorganisms can potentially be modified to increase the expression of pollutant-degrading enzymes, improve tolerance to toxic contaminants, or redirect metabolic pathways toward more complete degradation. Engineered microbial systems may be particularly useful for contaminants that are poorly degraded by naturally occurring microbial populations. However, environmental applications of genetically modified microorganisms require careful consideration of ecological stability, horizontal gene transfer, unintended interactions, containment, and regulatory requirements. For some applications, engineered enzymes may provide an alternative to releasing modified microorganisms into natural ecosystems.

Artificial intelligence and machine learning can complement these biological technologies by processing large datasets generated from environmental monitoring and omics analyses. Machine-learning models can identify relationships between microbial community composition, environmental parameters, pollutant concentrations, and remediation outcomes. In the future, predictive models may assist in selecting suitable microbial consortia, determining nutrient requirements, optimizing pH and oxygen conditions, and predicting treatment performance before large-scale implementation. Integration with automated monitoring systems could enable real-time adjustment of remediation conditions, transforming static treatment processes into adaptive systems.

Microbial electrochemical technologies are also receiving increasing attention because they connect microbial metabolism with controlled electron transfer. Electroactive microorganisms can transfer electrons to or from electrodes, allowing electrical conditions to influence microbial metabolic pathways. Microbial fuel cells, microbial electrolysis systems, and related technologies have been investigated for wastewater treatment, pollutant degradation, energy recovery, and resource recovery. Future research is likely to focus on improving electrode materials, biofilm stability, electron-transfer efficiency, reactor configuration, and overall energy balance. These technologies could become

particularly useful for contaminants whose degradation is limited by electron-donor or electron-acceptor availability.

Immobilized microbial cells and enzymes provide another promising direction. Immobilization can protect microorganisms or enzymes from unfavorable environmental conditions and facilitate their recovery and repeated use. Immobilized enzymes may be incorporated into treatment reactors, filtration systems, or engineered materials. Advances in nanotechnology and materials science may further improve enzyme stabilization and increase the surface area available for microbial attachment. Nevertheless, the environmental safety, production costs, regeneration capacity, and long-term stability of such materials require careful evaluation.

Future microbial remediation is also likely to emphasize resource recovery rather than contaminant removal alone. Wastewater and contaminated environments often contain valuable metals, nutrients, carbon compounds, and other resources. Microbial processes can potentially transform waste streams into useful products while simultaneously reducing environmental pollution. Metal-recovering microbial systems, nutrient recovery, microbial biomass production, bioenergy generation, and conversion of plastic-derived compounds into valuable chemicals represent examples of this circular approach. Such strategies can strengthen the economic and environmental sustainability of remediation technologies.

Field-scale implementation remains one of the major priorities for future research. Many microbial remediation studies are conducted under controlled laboratory conditions that do not reproduce the heterogeneity of natural environments. Future investigations should therefore combine laboratory experiments with pilot-scale and field-scale validation. Long-term monitoring should evaluate contaminant disappearance, transformation products, microbial community stability, toxicity reduction, ecosystem recovery, energy consumption, and greenhouse-gas emissions. Life-cycle assessment should be incorporated into technology development to determine whether an apparently biological solution provides genuine environmental benefits across its entire operational life.

Thus, the future of microbial remediation is likely to involve integrated systems rather than a single technology. Microbial communities, engineered enzymes, plants, electrochemical systems, sensors, computational models, and conventional treatment methods can potentially be combined to address complex contamination. Such integration can increase treatment flexibility while reducing dependence on intensive chemical or physical processes. The development of safe, scalable, economically feasible, and environmentally responsible technologies will be essential for translating advances in microbial biotechnology into practical remediation solutions.

V. INTEGRATED PHYTOMICROBIAL REMEDIATION FOR SUSTAINABLE ENVIRONMENTAL CLEANUP

Integrated phytomicrobial remediation combines the contaminant-removal capabilities of plants with the metabolic activities of microorganisms. Plants can absorb, stabilize, transform, or sequester certain contaminants, while microorganisms associated with plant roots can degrade organic pollutants, transform inorganic contaminants, and influence the availability and movement of chemicals in the rhizosphere. The interaction between plant roots and microorganisms creates a biologically active zone in soil known as the rhizosphere, where root-derived compounds, microbial metabolites, and environmental contaminants interact. This combined system provides an important approach for treating contaminated soils, sediments, and water while supporting ecological restoration.

Plants contribute to remediation through several mechanisms. Phytoextraction involves the uptake and accumulation of contaminants into plant tissues, whereas phytostabilization limits contaminant mobility by retaining pollutants within the soil or root zone. Phytodegradation involves transformation of contaminants within plant tissues, while rhizofiltration can remove contaminants from aqueous environments through plant-root interactions. However, the effectiveness of plants alone may be restricted by contaminant toxicity, low bioavailability, slow growth, limited root penetration, and the inability of plants to metabolize certain complex pollutants. Microorganisms can complement these limitations by transforming contaminants in the rhizosphere and modifying their chemical availability.

Root-associated microorganisms are particularly important because plants release a wide variety of organic compounds into the surrounding soil. Root exudates may include sugars, amino acids, organic acids, phenolic compounds, and other molecules that provide carbon and energy sources for microbial populations. These compounds can stimulate microbial growth and increase the abundance of microorganisms capable of degrading environmental contaminants. Consequently, plants can create favorable microhabitats for pollutant-degrading microorganisms while microorganisms can support plant development and stress tolerance.

Microbial communities associated with plant roots can degrade hydrocarbons, pesticides, dyes, pharmaceuticals, and other organic contaminants. Rhizosphere microorganisms may produce oxygenases, hydrolases, dehalogenases, laccases, peroxidases, and other enzymes involved in pollutant transformation. Some microorganisms can also produce biosurfactants that increase the apparent availability of hydrophobic pollutants, facilitating their microbial degradation. Others can modify soil pH, produce siderophores, mobilize nutrients, or alter metal speciation. These microbial functions can influence both contaminant fate and plant health.

The relationship between plants and microorganisms is often bidirectional. Plants provide microorganisms with nutrients and physical habitats, while beneficial microorganisms can improve plant growth by producing phytohormones, increasing nutrient availability, suppressing pathogens, and helping plants tolerate toxic environmental conditions. Plant growth-promoting microorganisms can therefore strengthen the capacity of vegetation to survive in contaminated environments. Improved plant growth leads to greater root development, which increases the volume of soil influenced by rhizosphere processes and can further stimulate microbial activity.

Integrated phytomicrobial systems can be particularly useful for large contaminated areas where intensive excavation or chemical treatment may be impractical. Vegetation-based approaches generally require less physical disturbance than excavation and can contribute to soil stabilization and landscape restoration. Establishment of vegetation can also reduce erosion and surface runoff, thereby limiting the movement of contaminants to surrounding ecosystems. However, the choice of plant species and microorganisms must be matched to the contaminant and environmental conditions. Plants should be capable of surviving at the contaminated site, while microbial populations should possess appropriate metabolic functions and ecological compatibility.

Omics technologies can improve understanding of phytomicrobial interactions by characterizing changes in microbial community composition, functional genes, gene expression, enzymes, and metabolites within the rhizosphere. Metagenomic and metatranscriptomic approaches can identify microorganisms and metabolic pathways enriched around plant roots. Metabolomics can further reveal interactions between root exudates, microbial metabolites, and pollutant transformation products. This information can help researchers identify plant-microorganism combinations that provide complementary remediation functions.

The integration of phytoremediation with microbial electrochemical technologies, bioaugmentation, biostimulation, and engineered microbial systems represents a further opportunity for future development. For example, plant-based remediation could provide ecological restoration while microbial systems target specific contaminants that are poorly removed by plants alone. Similarly, targeted nutrient management could stimulate naturally occurring rhizosphere microorganisms without requiring large-scale introduction of external microbial strains.

VI. CONCLUSION

Microbial metabolism provides a versatile biological platform for addressing diverse environmental contamination problems. The ability of microorganisms to oxidize, reduce, hydrolyze, dehalogenate, immobilize, adsorb, transform, and mineralize pollutants provides multiple routes for remediation. Bacteria, fungi, archaea, algae, and microbial consortia can participate in these processes through highly diverse metabolic pathways and enzyme systems.

The potential of microbial remediation extends across petroleum hydrocarbons, pesticides, pharmaceuticals, dyes, heavy metals, plastics, and complex wastewater contaminants. For organic pollutants, microbial metabolism can convert complex compounds into simpler intermediates and, under appropriate conditions, ultimately into carbon dioxide, water, biomass, and inorganic products. For metals, microorganisms can reduce mobility and bioavailability through biosorption, bioaccumulation, precipitation, and redox transformation.

Emerging technologies are significantly expanding this potential. Microbial consortia can provide complementary metabolic capabilities, while metabolic engineering and synthetic biology can increase pathway specificity. Omics technologies enable researchers to characterize microbial functions at unprecedented resolution. Enzyme engineering and immobilization offer controlled alternatives to whole-cell remediation. Microbial electrochemical technologies provide additional possibilities for manipulating electron transfer, treating pollutants, recovering resources, and potentially producing energy.

Nevertheless, microbial remediation should not be considered a universal replacement for physicochemical technologies. Its effectiveness depends strongly on environmental conditions, pollutant bioavailability, microbial ecology, nutrient availability, toxicity, and treatment objectives. Careful assessment of transformation products and ecological consequences is essential. Hybrid approaches combining biological, chemical, physical, electrochemical, and phytoremediation technologies may therefore be particularly useful for complex contamination.

The future of sustainable environmental remediation lies in understanding and controlling microbial metabolism rather than simply introducing microorganisms into contaminated environments. Integrating microbial ecology with omics, metabolic engineering, artificial intelligence, biosensors, advanced reactor design, and resource recovery can create more targeted and adaptable remediation systems. The development of such systems can help transform environmental biotechnology from a contaminant-removal discipline into a broader platform for ecosystem restoration, resource recovery, and circular environmental management.

REFERENCES

- [1]. Kuppan, N., Padman, M., Mahadeva, M., Srinivasan, S., & Devarajan, R. (2024). A comprehensive review of sustainable bioremediation techniques: Eco friendly solutions for waste and pollution management. *Waste Management Bulletin*, 2(3), 154–171. <https://doi.org/10.1016/J.WMB.2024.07.005> (DOI)
- [2]. Monga, D., & Kaur, P. (2022). Microbe mediated remediation of dyes, explosive waste and polyaromatic hydrocarbons, pesticides and pharmaceuticals. *Environmental Technology & Innovation*. (PubMed Central (PMC))
- [3]. Sreedevi, P. R., Suresh, K., & Jiang, G. (2022). Bacterial bioremediation of heavy metals in

- wastewater: A review of processes and applications. *Journal of Water Process Engineering*, 48, 102884. <https://doi.org/10.1016/j.jwpe.2022.102884> (ScienceDirect)
- [4]. Tufail, M. A., Iltaf, J., Zaheer, T., et al. (2022). Recent advances in bioremediation of heavy metals and persistent organic pollutants: A review. *Science of the Total Environment*, 850, 157961. <https://doi.org/10.1016/j.scitotenv.2022.157961> (ScienceDirect)
- [5]. Sikarwar, D., & Bashir, Y. (2024). Critical perspective on the elimination of emerging contaminants from industrial wastewater via microbial electrochemical technologies. *CLEAN – Soil, Air, Water*. <https://doi.org/10.1002/clen.202300346> (Wiley Online Library)
- [6]. Pandolfo, E., Barra Caracciolo, A., & Rolando, L. (2023). Recent advances in bacterial degradation of hydrocarbons. *Water*, 15(2), 375. <https://doi.org/10.3390/w15020375> (MDPI)
- [7]. Varjani, S. J. (2017). Microbial degradation of petroleum hydrocarbons. *Bioresource Technology*, 223, 277–286. (ScienceDirect)
- [8]. Lan, F., Ma, H., Zhang, X., Liu, C., & Chen, J. (2024). A review of research progress of current status, influence factors, degradation of typical pollutants by microbial electrolytic cell. *Journal of Water Process Engineering*, 60, 106452. <https://doi.org/10.1016/j.jwpe.2024.106452> (ScienceDirect)
- [9]. Wang, X., Aulenta, F., Puig, S., Esteve-Núñez, A., He, Y., Mu, Y., & Rabaey, K. (2020). Microbial electrochemistry for bioremediation. *Environmental Science and Ecotechnology*, 1, 100013. <https://doi.org/10.1016/j.es.2020.100013> (PubMed Central (PMC))
- [10]. Schneier, A., Melaugh, G., & Sadler, J. C. (2024). Engineered plastic-associated bacteria for biodegradation and bioremediation. *Biotechnology for the Environment*, 1. (Springer)
- [11]. Dhali, S. L., Parida, D., Kumar, B., & Bala, K. (2024). Recent trends in microbial and enzymatic plastic degradation: A solution for plastic pollution predicaments. *Biotechnology for Sustainable Materials*, 1, 11. (Springer)
- [12]. Sikarwar, D., & Bashir, Y. (2024). Microbial electrochemical technologies for emerging contaminant remediation. *CLEAN – Soil, Air, Water*. (Wiley Online Library)
- [13]. A comprehensive review on the potential of microbial enzymes in multipollutant bioremediation: Mechanisms, challenges, and future prospects. (2023). *Journal of Environmental Management*, 334, 117532. <https://doi.org/10.1016/j.jenvman.2023.117532> (ScienceDirect)
- [14]. Microbial plastic degradation: Enzymes, pathways, challenges, and perspectives. (2025). *Microbiology and Molecular Biology Reviews*. (ASM Journals)
- [15]. Riseh, R. S., Vazvani, M. G., Hajabdollahi, N., et al. (2023). Bioremediation of heavy metals by rhizobacteria. *Applied Biochemistry and Biotechnology*, 195, 4689–4711. (Springer)
- [16]. A review on catalytic-enzyme degradation of toxic environmental pollutants: Microbial enzymes. (2021). *Environmental Technology & Innovation*. (PubMed)
- [17]. Recent advances in microbial and enzymatic engineering for the biodegradation of micro- and nanoplastics. (2024). *RSC Advances*, 14, 9943–9966. <https://doi.org/10.1039/D4RA00844H> (ScienceDirect)